

Pembrolizumab Paediatric Monotherapy

INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
As monotherapy for the treatment of paediatric patients aged 3 years and older with relapsed or refractory (R/R) classical Hodgkin lymphoma (cHL) who have failed autologous stem cell transplant (ASCT) or following at least two prior therapies when ASCT is not a treatment option.	C81	P00711a	ODMS 01/02/2022

TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patients individual clinical circumstances.

Pembrolizumab is administered once every 21 days until disease progression or unacceptable toxicity.

Atypical responses (i.e. an initial transient increase in tumour size or small new lesions within the first few months followed by tumour shrinkage) have been observed. It is recommended to continue treatment for clinically stable patients with initial evidence of disease progression until disease progression is confirmed.

Facilities to treat anaphylaxis MUST be present when systemic anti-cancer therapy (SACT) is administered.

Day	Drug	Dose	Route	Diluent & Rate	Cycle
1	Pembrolizumab	2mg/kg (up to a maximum of 200mg)	IV infusion	50ml 0.9% NaCl over 30 minutes	Every 21 days
Pembrolizumab is diluted to a final concentration ranging from 1-10mg/ml.					
Administer using a low-protein binding 0.2 to 5 micrometre in-line or add-on filter.					

ELIGIBILITY:

- Indication as above
- ECOG status 0-1
- Adequate haematological, hepatic and renal function

CAUTION:

- History of serious autoimmune disease

EXCLUSIONS:

- Known hypersensitivity to pembrolizumab or to any of the excipients
- Known clinically active central nervous system (CNS) involvement
- Active autoimmune disease that has required systemic treatment in the past 2 years
- Information regarding prior therapy with an anti PD-1 or anti PD-L1 antibody is available [here](#)

NCCP Regimen: Pembrolizumab Paediatric Therapy	Published: 1/12/2023 Review: 1/12/2024	Version number: 1
Tumour Group: Lymphoma NCCP Regimen Code: P00711a	IHS Contributor: Dr Jane Pears	Page 1 of 5

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

- Any medical condition that requires immunosuppressive doses of systemic corticosteroids or other immunosuppressive medication(s) (defined as >10mg prednisolone/daily (or steroid equivalent, excluding inhaled or topical steroids)
- History of interstitial lung disease
- Any active clinically significant infection requiring therapy
- Pregnancy and breastfeeding

PRESCRIPTIVE AUTHORITY:

The treatment plan must be initiated by a Consultant Medical Oncologist or by a Consultant Haematologist working in the area of haematological malignancies.

TESTS:

Baseline tests:

- FBC, renal and liver profile
- Glucose
- Thyroid function tests
- Virology Screen: Hepatitis B (HBsAg, HBcoreAb) and Hepatitis C

Regular tests:

- FBC, renal and liver profile prior to each cycle
- Glucose prior to each cycle
- Thyroid function tests every 3 to 6 weeks

Disease monitoring:

Disease monitoring should be in line with the patient’s treatment plan and any other test/s as directed by the supervising Consultant.

DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant.
- Management of immune-related adverse reactions may require withholding of a dose or permanent discontinuation of pembrolizumab therapy and institution of systemic high-dose corticosteroid.
- Dose reduction is not recommended.
- Guidelines for withholding of doses or permanent discontinuation are described below in Table 1.

NCCP Regimen: Pembrolizumab Paediatric Therapy	Published: 1/12/2023 Review: 1/12/2024	Version number: 1
Tumour Group: Lymphoma NCCP Regimen Code: P00711a	IHS Contributor: Dr Jane Pears	Page 2 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

Table 1: Recommended treatment modifications for pembrolizumab

Immune-related adverse reactions	Severity (NCI-CTCAE v.4 grading)	Treatment modification
Pneumonitis	Grade 2	Withhold*
	Grade 3 or 4, or recurrent Grade 2	Permanently discontinue
Colitis	Grade 2 or 3	Withhold*
	Grade 4 or recurrent Grade 3	Permanently discontinue
Nephritis	Grade 2 with creatinine > 1.5 to ≤ 3 times upper limit of normal (ULN)	Withhold*
	Grade ≥ 3 with creatinine > 3 times ULN	Permanently discontinue
Endocrinopathies	Grade 2 adrenal insufficiency and Hypophysitis	Withhold treatment until controlled by hormone replacement
	Grades 3 or 4 adrenal insufficiency or symptomatic hypophysitis	Withhold*
	Type 1 diabetes associated with Grade ≥ 3 hyperglycaemia (glucose > 250 mg/dL or > 13.9 mmol/L) or associated with ketoacidosis	For patients with Grade 3 or Grade 4 endocrinopathy that improved to Grade 2 or lower and is controlled with hormone replacement, if indicated, continuation of pembrolizumab may be considered after corticosteroid taper, if needed. Otherwise treatment should be discontinued.
	Hyperthyroidism Grade ≥ 3	
Hypothyroidism	Hypothyroidism may be managed with replacement therapy without treatment interruption.	
Hepatitis	Grade 2 with aspartate aminotransferase (AST) or alanine aminotransferase (ALT) > 3 to 5 times ULN or total bilirubin > 1.5 to 3 times ULN	Withhold*
	Grade ≥ 3 with AST or ALT > 5 times ULN or total bilirubin > 3 times ULN	Permanently discontinue
	In case of liver metastasis with baseline Grade 2 elevation of AST or ALT, hepatitis with AST or ALT increases ≥ 50% and lasts ≥ 1 week	
Skin reactions	Grade 3 or suspected Stevens-Johnson syndrome (SJS) or toxic epidermal necrolysis (TEN)	Withhold*
	Grade 4 or confirmed SJS or TEN	Permanently discontinue
Other immune-related adverse reactions**	Based on severity and type of reaction (grade 2 or Grade 3)	Withhold*
	Grade 3 or 4 myocarditis	Permanently discontinue
	Grade 3 or 4 encephalitis	
	Grade 3 or 4 Guillain-Barre syndrome	
Grade 4 or recurrent Grade 3		
Infusion-related reactions	Grade 3 or 4	Permanently discontinue

*Until adverse reactions recover to Grade 0-1. If treatment related toxicity does not resolve to Grade 0-1 within 12 weeks after last dose of pembrolizumab or if corticosteroid dosing cannot be reduced to ≤ 10mg prednisone or equivalent per day within 12 weeks, pembrolizumab should be permanently discontinued

NCCP Regimen: Pembrolizumab Paediatric Therapy	Published: 1/12/2023 Review: 1/12/2024	Version number: 1
Tumour Group: Lymphoma NCCP Regimen Code: P00711a	IHS Contributor: Dr Jane Pears	Page 3 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

**Pembrolizumab should be permanently discontinued for Grade 4 or recurrent Grade 3 immune-related adverse reactions, unless otherwise specified in Table 1.

Renal and Hepatic Impairment:

Table 2: Dose modification of pembrolizumab in renal and hepatic impairment

Renal Impairment		Hepatic Impairment	
Mild/Moderate	No dose adjustment required	Mild	No dose adjustment required
Severe	Has not been studied	Moderate/Severe	Has not been studied

SUPPORTIVE CARE:

EMETOGENIC POTENTIAL: Minimal (Refer to local policy)

PREMEDICATIONS: Not usually required

OTHER SUPPORTIVE CARE: Not usually required

ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS:

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

- Immune-mediated adverse reactions:** Most immune-related adverse reactions occurring during treatment with pembrolizumab are reversible and managed with interruptions of pembrolizumab, administration of corticosteroids and/or supportive care. Immune-related adverse reactions have also occurred after the last dose of pembrolizumab. For suspected immune-related adverse reactions, adequate evaluation to confirm aetiology or exclude other causes should be ensured. Based on the severity of the adverse reaction, pembrolizumab should be withheld and corticosteroids administered. Upon improvement to Grade ≤ 1 , corticosteroid taper should be initiated and continued over at least 1 month.

Based on limited data from clinical studies in patients whose immune-related adverse reactions could not be controlled with corticosteroid use, administration of other systemic immunosuppressants can be considered. Pembrolizumab may be restarted within 12 weeks after last dose of pembrolizumab if the adverse reaction remains at Grade ≤ 1 and corticosteroid dose has been reduced to ≤ 10 mg prednisone or equivalent per day. Pembrolizumab must be permanently discontinued for any Grade 3 immune-related adverse reaction that recurs and for any Grade 4 immune-related adverse reaction toxicity, except for endocrinopathies that are controlled with replacement hormones. Specific guidelines for management of Immune Mediated Adverse Events are available.
- Infusion-related reactions:** Severe infusion-related reactions have been reported in patients receiving pembrolizumab. For severe infusion reactions, infusion should be stopped and pembrolizumab permanently discontinued. Patients with mild or moderate infusion reaction may continue to receive pembrolizumab with close monitoring; premedication with antipyretic and antihistamine may be considered.

NCCP Regimen: Pembrolizumab Paediatric Therapy	Published: 1/12/2023 Review: 1/12/2024	Version number: 1
Tumour Group: Lymphoma NCCP Regimen Code: P00711a	IHS Contributor: Dr Jane Pears	Page 4 of 5

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

DRUG INTERACTIONS:

- No formal pharmacokinetic drug interaction studies have been conducted with pembrolizumab. Since pembrolizumab is cleared from the circulation through catabolism, no metabolic drug-drug interactions are expected.
- The use of systemic corticosteroids or immunosuppressants before starting pembrolizumab should be avoided because of their potential interference with the pharmacodynamic activity and efficacy of pembrolizumab. However, systemic corticosteroids or other immunosuppressant's can be used after starting pembrolizumab to treat immune-related adverse reactions.
- Current drug interaction databases should be consulted for more information.

COMPANY SUPPORT RESOURCES/Useful Links:

Please note that this is for information only and does not constitute endorsement by the NCCP

Patient Guide

<https://www.hpra.ie/img/uploaded/swedocuments/196f9071-00a4-4498-9dcb-e29ef7b35e55.pdf>

Patient Alert Card

<https://www.hpra.ie/img/uploaded/swedocuments/c0984994-f8e8-4b10-95dd-7be12ff6c6f9.pdf>

REFERENCES:

1. Georger B et al. Pembrolizumab in paediatric patients with advanced melanoma or a PD-L1-positive, advanced, relapsed, or refractory solid tumour or lymphoma (KEYNOTE-051): interim analysis of an open-label, single-arm, phase 1-2 trial. *Lancet Oncol.* 2020 Jan; 21(1):121-133. doi: 10.1016/S1470-2045(19)30671-0. Epub 2019 Dec 4. PMID: 31812554.
2. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V5 2023. Available at: <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp-classification-document-for-systemic-anti-cancer-therapy-sact-induced-nausea-and-vomiting.pdf>
3. Pembrolizumab (Keytruda®). Summary of Product Characteristics. Last updated: 17/11/2022. Accessed November 2023. Available at: https://www.ema.europa.eu/en/documents/product-information/keytruda-epar-product-information_en.pdf

Version	Date	Amendment	Approved By
1	1/12/2023		Dr Jane Pears

Comments and feedback welcome at oncologydrugs@cancercontrol.ie.

NCCP Regimen: Pembrolizumab Paediatric Therapy	Published: 1/12/2023 Review: 1/12/2024	Version number: 1
Tumour Group: Lymphoma NCCP Regimen Code: P00711a	IHS Contributor: Dr Jane Pears	Page 5 of 5

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens