

Drug	Dose	Duration	+/- Notes
Uncomplicated <u>anogenital and pharyngeal</u> gonorrhoea in Adults			
Ceftriaxone	1gram deep IM stat	Stat dose	Dissolve 1g ceftriaxone in 3.5ml of 1% Lidocaine Injection for IM injection. Not for IV injection
Uncomplicated <u>anogenital</u> gonorrhoea in Adults with cephalosporin allergy* OR previous or immediate and/or severe hypersensitivity to penicillin or other β-lactam*. Refer to dedicated GUM service and where not possible discuss with a specialist in GUM or ID			
Ciprofloxacin	500mg PO stat	Stat dose	When known to be sensitive to quinolones Avoid ciprofloxacin in pregnancy. Caution with use of ciprofloxacin- see Fluoroquinolone warning below.
OR			
Azithromycin	2g PO stat	Stat dose	When known to be susceptible to Azithromycin Tablets: Take with or without food Capsules: Take 1 hour before or 2 hours after food
Pharyngeal gonorrhoea⁺ in Adults with cephalosporin allergy* OR previous or immediate and/or severe hypersensitivity to penicillin or other β-lactam* Refer to dedicated GUM service and where not possible discuss with a specialist in GUM or ID			
Ciprofloxacin	500mg PO every 12 hours x 2 doses	1 day	When known to be sensitive to quinolones Avoid ciprofloxacin in pregnancy. Caution with use of ciprofloxacin- see Fluoroquinolone warning below.
OR			
Azithromycin	2g PO stat	Stat dose	When known to be susceptible to Azithromycin Tablets: Take with or without food Capsules: Take 1 hour before or 2 hours after food